



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A23L 2/66</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/39641</b> <b>(43) International Publication Date:</b> 30 October 1997 (30.10.97)
<b>(21) International Application Number:</b> PCT/DK97/00177 <b>(22) International Filing Date:</b> 18 April 1997 (18.04.97) <b>(30) Priority Data:</b> 0457/96                      18 April 1996 (18.04.96)                      DK <b>(71) Applicant (for all designated States except US):</b> M.D. FOODS A.M.B.A. [DK/DK]; Skanderborgvej 277, DK-8260 Viby J. (DK). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BERTHELSEN, Jørn, Havskov [DK/DK]; Lappestensvej 14B, DK-3000 Helsingør (DK). FRØKJÆR, Sven [DK/DK]; Solvej 6, DK-2840 Holte (DK). SALTIN, Bengt [SE/DK]; Vindekilde Strandvej 30, DK-4540 Faarevejle (DK). <b>(74) Agent:</b> HOFMAN-BANG & BOUTARD, LEHMANN & REE A/S; Hans Bekkevolds Allé 7, DK-2900 Hellerup (DK).		<b>(81) Designated States:</b> AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ADDITIVE FOR USE IN ANY ENERGY SUPPLEMENTATION, USE OF A PROTEIN HYDROLYSATE TO THE PREPARATION OF ENERGY SUPPLEMENTATION AND AN ENERGY SUPPLEMENT CONTAINING SUCH AN ADDITIVE  <b>(57) Abstract</b>  Additive for use in any energy supplementation or metabolic nutrient in the form of a beverage or other nutrient for athletes or other persons in need of increased glycogen level, use of a protein hydrolysate to the preparation of such an energy supplementation and an energy supplement containing such an additive. The protein hydrolysate can be of animal or vegetable origin.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Additive for use in any energy supplementation, use of a protein hydrolysate to the preparation of energy supplementation and an energy supplement containing such an additive

- 5 The present invention relates to an additive for use in any energy supplementation or metabolic nutrient, the use of a protein hydrolysate to the preparation of an energy supplementation or metabolic nutrient. The energy supplementation of the invention can be in the form of a  
10 beverage or other nutrient preparation for an athlete or another person in need of increased glycogen level and an additive for such a beverage or other nutrient for oral intake.

- Athletes consume energy, primarily glycogen  
15 (carbohydrate) and later on (after 40 minutes) fat. Carbohydrate is a fast and light energy source. Consequently, carbohydrates are taken in together with liquid. In order that carbohydrates can be utilised they must pass from the blood into muscles. This transport  
20 requires insulin.

As it is, intake of water + carbohydrate increases insulin secretion moderately.

- According to the invention a study has surprisingly shown that the combination water + carbohydrates + special  
25 protein hydrolysate increases insulin secretion dramatically.

- This is interesting because insulin is considered an anabolic (building) hormone, and is necessary both for optimum energy uptake in muscles and brain and for uptake  
30 in (transport to) the muscles of amino acids, and is also necessary when rebuilding the spent glycogen.

This means that

1) the special protein hydrolysate of the invention + carbohydrate + water increases insulin secretion and consequently the chance of supplying more energy to  
5 muscles when exercising than carbohydrate + water or carbohydrate + water + intact protein.

2) after the metabolic stress, intake of the energy supplement of the invention in the form of a beverage or other nutrient entails faster rebuilding of glycogen  
10 depots and faster rebuilding of degraded muscular proteins.

The additive according to the invention is characterized in that it is a protein hydrolysate having a degree of hydrolysis (DH) of 1-50, preferably 15-30 and especially  
15 about 25, which protein hydrolysate after oral intake rises the insulin secretion to a level above what is achieved by intake of the intact parent protein. Preferably the protein hydrolysate does not increase the gastric emptying, generally gives persons less satiety,  
20 increases the glycogen synthesis or rebuilding after depletion and reduces muscle break-down during metabolic stress, due to increased influx of amino acids into the muscle compared to what is achieved by intake of the intact parent protein.

25 Studies have surprisingly shown that both animal and vegetable protein is useful in the invention, which consequently comprises both types.

The protein hydrolysate can thus be of animal or vegetable origin. Useful protein hydrolysates are whey  
30 protein hydrolysates, casein protein hydrolysates, soy protein hydrolysates, wheat protein hydrolysates or pea protein hydrolysates.

The use of a protein hydrolysate to the preparation of an energy supplement or metabolic nutrient is as defined in claims 7 to 12 and the energy supplementation as defined in the claims 13 to 19.

- 5 The proteins useful in the invention can be obtained by enzymatic or non-enzymatic hydrolysis e.g. by enzymatic hydrolysis as described in the international patent application WO 92/21248.

- 10 An additive according to the invention can be used in a beverage or other nutrient product for athletes to re-establish liquid and salt balance as well as glycogen levels in the body before, under or after sport or other physical work. The beverage is preferable an isotonic fluid. The nutrient can be liquid as well as solid or be  
15 a mixture of liquids and solids, such as a paste. It can also contain sweeteners, flavours and other ingredients normally contained in beverages of this kind.

- The energy supplementation of the invention contains a protein hydrolysate, a carbohydrate and optionally water  
20 or it is taken together with water.

The carbohydrate can be glucose, sucrose, maltose or maltodextrines or mixtures thereof.

- The following is an example of a useful energy supplementation in the form of a beverage according to  
25 the invention.

**GENERAL EXAMPLE**

Ingredients, pr. liter:

Protein hydrolysate	30 g ± 15 g
Carbohydrate	70 g ± 30 g
Sodium	25 mM ± 10 mM
Potassium	3 mM ± 2 mM
Chloride	12 mM ± 6 mM

Carbohydrate can be glucose, sucrose, maltose and/or  
5 maltodextrines.

Trace amounts of other mineral can be present.

**SPECIFIC EXAMPLE**

	Lacprodan DI-3065	3.75%
	SPG 30 (DE 28-32)	3.50%
10	SPG 20 (DE 20-23)	3.50%
	NaCl	0.07%
	NaOH	0.03%
	Aspartam	0.03%
	Flavour	0.40%
15	Malic Acid	0.45%
	Water	88.27%

**EXPERIMENTS**

The following experiments have been carried out.

- 20 RATE OF GASTRIC EMPTYING AND PLASMA INSULIN RESPONSE TO  
PROTEIN HYDROLYSATE SOLUTIONS.

The purpose of the present study was to determine the rate of gastric emptying of four solutions containing different quantities of proteins and protein hydrolysates (approx. 4 amino acid residuals).

- 5 Six healthy human subjects (three males and three females;  $22 \pm 2.1$  years;  $73.2 \pm 7.8$  kg weight and  $177.3 \pm 10.00$  cm height) participated in this study, which was approved by the local ethics committee. Test meals (600 ml) were administered through a nasogastric tube, 30 minutes after
- 10 a gastric washout. Four different meals were studied in random order, over a period of 4 weeks. Solution A (control) was composed of glucose ( $0.025 \text{ g ml}^{-1}$ ) and NaCl (0.9%). The three other solutions contained similar quantities of protein ( $0.25 \text{ g (kg body mass)}^{-1}$ ) and
- 15 glucose ( $0.025 \text{ g ml}^{-1}$ ). Solutions were composed of: B, a pea protein hydrolysate and NaCl; C, a whey protein hydrolysate and NaCl; and D, milk powder. Solutions B and C were matched for osmolality, caloric density, and nitrogen contents, but contained different amounts of
- 20 essential amino acids (35.5 and 45.1%, respectively). and non-essential amino acids (64.5 and 54.9%, respectively). Each meal was delivered at  $37^\circ\text{C}$ , after adjusting it to a pH of approx. 7. During the 3 h that each trial lasted, blood samples were obtained every 20 min. for the
- 25 assessment of the plasmatic concentrations of glucose and insulin. Gastric samples were taken every 10 min, to determine the volume remaining in the stomach using George's double-sampling technique, as applied by Beekers et al. (1988).
- 30 The rate of gastric emptying followed an exponential pattern in all cases ( $r = 0.86\text{--}0.99$ ). Solution A was emptied faster than the other three solutions with a half-time of  $9.4 \pm 1.2$  min ( $p < 0.05$ ). However, D emptied at a slower rate ( $26.4 \pm 10.0$  min,  $p < 0.05$ , paired t test) than

both B ( $16.3 \pm 5.4$  min) and C ( $17.2 \pm 6.1$  min). Despite important differences in amino acid composition, both protein hydrolysate solutions emptied at a similar rate. Caloric density was closely related to the half-time of gastric emptying ( $r = 0.96$ ,  $p < 0.05$ ,  $n = 4$ ) and with the rate at which calories were delivered to the duodenum ( $r = 0.99$ ,  $p < 0.001$ ,  $n = 4$ ). The insulin response was significantly different among solutions ( $p < 0.001$ , ANOVA). The plasma insulin peak elicited by the C and B ( $80.3 \pm 50.7$  and  $84.8 \pm 34.9$ ,  $\mu\text{mol ml}^{-1}$ , respectively) were approx. 2- and 4-fold greater than that produced by D ( $47.8 \pm 13.7$   $\mu\text{mol ml}^{-1}$ ,  $p < 0.05$ , paired t test) and A ( $28.7 \pm 18.0$   $\mu\text{mol ml}^{-1}$   $p < 0.05$ , paired t test).

The results of the present study suggest that protein hydrolysate solutions differing in the amino acid composition, but matched for temperature, volume, osmolality, acidity and caloric density, are emptied at a similar rate. In addition, the ingestion of solutions combining glucose and protein hydrolysates produce a significant and unexpected increase in plasma insulin concentration.



## PATENT CLAIMS

1. Additive for use in any energy supplementation or metabolic nutrient, c h a r a c t e r i z e d by the fact that it is a protein hydrolysate having a degree of hydrolysis (DH) of 1-50, which protein hydrolysate after oral intake rises the insulin secretion to a level above what is achieved by intake of the intact parent protein.
2. Additive according to claim 1, c h a r a c t e r i z e d by the fact that it is a protein hydrolysate which does not delay the gastric emptying, generally gives persons less satiety, optimizes the glycogen synthesis after depletion and reduces muscle break-down during metabolic stress, due to increased influx of amino acids into the muscle compared with what is achieved by intake of the intact parent protein.
3. Additive according to claim 1 or 2, c h a r a c t e r i z e d by the fact that it is a protein hydrolysate having a degree of hydrolysis of 15 - 30.
4. Additive according to claim 1 or 2, c h a r a c t e r i z e d by the fact that it is a protein hydrolysate having a degree of hydrolysis of about 25.
5. Additive according to any of the preceding claims, c h a r a c t e r i z e d in, that it is an animal or vegetable protein hydrolysate.
6. Additive according to claim 5, c h a r a c t e r i z e d in that it is an animal or vegetable protein hydrolysate.
7. Use of a protein hydrolysate to the preparation of a energy supplementation or metabolic nutrient, c h a r a c t e r i z e d in that it is a protein hydrolysate having a degree of hydrolysis (DH) of 1-50, which protein

hydrolysate after oral intake rises the insulin secretion to a level above what is achieved by intake of the intact parent protein.

8. Use of a protein hydrolysate to the preparation of a  
5 energy supplementation or metabolic nutrient, c h a -  
r a c t e r i z e d in that it is a protein hydrolysate  
which does not delay the gastric emptying, generally  
gives persons less satiety, optimizes the glycogen  
10 synthesis after depletion and reduces muscle break-down  
during metabolic stress, due to increased influx of amino  
acids into the muscle compared with what is achieved by  
intake of the intact parent protein.

9. Use of a protein hydrolysate to the preparation of a  
energy supplementation or metabolic nutrient, c h a -  
15 r a c t e r i z e d by the fact that it is a protein  
hydrolysate having a degree of hydrolysis of 15 - 30.

10. Use of a protein hydrolysate to the preparation of a  
energy supplementation or metabolic nutrient, c h a -  
r a c t e r i z e d by the fact that it is a protein  
20 hydrolysate having a degree of hydrolysis of about 25.

11. Use of a protein hydrolysate to the preparation of a  
energy supplementation or metabolic nutrient, c h a -  
r a c t e r i z e d in, that it is an animal or vegetable  
protein hydrolysate.

25 12. Use of a protein hydrolysate to the preparation of a  
energy supplementation or metabolic nutrient, c h a -  
r a c t e r i z e d in that it is a whey protein  
hydrolysate, a casein protein hydrolysate, a soy protein  
hydrolysate, a wheat protein hydrolysate or a pea protein  
30 hydrolysate.

13. Energy supplementation in the form of a beverage or other nutrient for human beings characterized in, that it contains an additive according to any of the claims 1 - 6 and a carbohydrate.

5 14. Energy supplementation according to claim 13, characterized in that the additive is a protein hydrolysate having a degree of hydrolysis (DH) of 1-50, which protein hydrolysate after oral intake rises the insulin secretion to a level above what is achieved by  
10 intake of the intact parent protein.

15 15. Energy supplementation according to claim 13, characterized in that the additive is a protein hydrolysate which does not delay the gastric emptying, generally gives persons less satiety, optimizes the glycogen synthesis after depletion and reduces muscle break-down during metabolic stress, due to increased influx of amino acids into the muscle compared with what is achieved by intake of the intact parent protein.

20 16. Energy supplementation according to claim 13, characterized in that the additive is a protein hydrolysate having a degree of hydrolysis of 15 - 30.

25 17. Energy supplementation according to claim 13, characterized in that the additive is a protein hydrolysate having a degree of hydrolysis about 25.

18. Energy supplementation according to claim 13, characterized in that the additive is an animal or vegetable protein hydrolysate.

30 19. Energy supplementation according to claim 13, characterized in that the additive is a whey protein hydrolysate, a casein protein hydrolysate, a soy

protein hydrolysate, a wheat protein hydrolysate or a pea protein hydrolysate.

20. Energy supplementation according to any of the claims 13 to 19, characterized in that the
- 5 carbohydrate is glucose, sucrose, maltose or a maltodextrine or mixtures thereof.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00177

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A23L 2/66

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, DIALINDEX (FOODSCIENCE)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9215696 A1 (NOVO NORDISK A/S), 17 Sept 1992 (17.09.92), page 1, line 23 - line 25; page 2, line 1 - line 3; page 4, line 8 - line 16, page 15, line 12 - line 16, abstract --	1-20
X	WO 9215697 A1 (NOVO NORDISK A/S), 17 Sept 1992 (17.09.92), page 2, line 11 - line 13, claims 1, 16, examples 5,6, abstract --	1-20
A	US 4107334 A (RAMESH C. JOLLY), 15 August 1978 (15.08.78), claims 1,8 -----	1,13,20

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 July 1997

Date of mailing of the international search report

24-07-1997

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Sofia Nikolopoulou  
Telephone No. +46 8 782 25 00

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

01/07/97

International application No.  
PCT/DK 97/00177

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9215696 A1	17/09/92	AU 650524 B	23/06/94
		AU 1428192 A	06/10/92
		CA 2105673 A	08/09/92
		EP 0575452 A	29/12/93
		JP 6504677 T	02/06/94
		NO 933163 D	00/00/00
-----			
WO 9215697 A1	17/09/92	AU 1422492 A	06/10/92
		CA 2104595 A	08/09/92
		EP 0575443 A,B	29/12/93
		JP 6505162 T	16/06/94
		US 5520935 A	28/05/96
-----			
US 4107334 A	15/08/78	BE 859635 A	12/04/78
		CA 1094867 A	03/02/81
		DE 2745954 A,C	20/04/78
		FR 2367773 A,B	12/05/78
		GB 1536990 A	29/12/78
		JP 53047560 A	28/04/78
		LU 78294 A	01/06/79
		NL 7710841 A	17/04/78
SE 7710423 A	14/04/78		
-----			